



Dr. Scholl's® Duragel® Callus Remover

The Drug labeling information on this website may differ from the product labeling on purchased product. Please consult your purchased package for information specific to that product.

Drug Facts	
Active ingredient Salicylic acid 40%.....	PurposeCallus remover
Uses <ul style="list-style-type: none"> ▪ for the removal of calluses ▪ relieves pain by removing calluses 	
Warnings For external use only	
Do not use <ul style="list-style-type: none"> ▪ if you are a diabetic ▪ if you have poor blood circulation ▪ on irritated skin or any area that is infected or reddened 	
Stop use and ask a doctor if discomfort lasts	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions <ul style="list-style-type: none"> ▪ wash affected area and dry thoroughly ▪ if necessary, cut medicated disc to fit callus ▪ apply medicated disc with sticky side adhering to skin ▪ cover medicated disc with enclosed cushion ▪ after 48 hours, remove medicated disc ▪ repeat procedure every 48 hours as needed for up to 14 days (until callus is removed) ▪ may soak callus in warm water for 5 minutes to assist in removal 	
Other information <ul style="list-style-type: none"> ▪ Store between 20° to 25°C (68° to 77°F). ▪ Protect from excessive heat (temperatures above 40°C (104°F)). 	
Inactive ingredients antioxidant (CAS 991-84-4), iron oxides, mineral oil, petroleum hydrocarbon resin, silicon dioxide, synthetic polyisoprene rubber, talc	
Questions? 1- 866-360-3226	